# **WEST Search History**

Hide Items Restore Clear Cancel

DATE: Thursday, June 23, 2005

Hide?	<u>Set</u> Name	Query	<u>Hit</u> Count
	DB=I	PGPB, USPT, USOC, EPAB, JPAB, DWPI; PLUR = YES; OP = OR	
Γ	L17	(glutamine near derivative) and ((gamma adj2 glutamyltranferase) or (alpha adj2 glutamyltraspeptidase) or (gamma adj2 glutamyl adj (peptidyltransferase or transpeptidase)) or (gamma adj glutamyltransferase) or (gamma adj gpt) or ??gpt or (gamma adj gt) or ??gt or (gamma adj gtp) or ??gtp)	ç
Γ	L16	(glutamine near derivative) and ((diabetic adj neuropath\$) or glomeruloscleros\$ or glomerulopath\$ or ((chronic or persistent) adj (kidney or renal) adj disease))	5
Γ	L15	ll1 and ((gamma adj2 glutamyltranferase) or \$glutamyltransferase or (alpha adj2 glutamyltraspeptidase) or \$glutamyltranspeptidase or (gamma adj2 glutamyl adj (peptidyltransferase or transpeptidase)) or (\$glutamyl adj (peptidyltranferase or transpeptidase)) or (gamma adj glutamyltransferase) or \$glutamyltrasferase or (gamma adj gpt) or \$gpt or (gamma adj gt) or \$gt or (gamma adj gtp) or \$gt)	333
Γ	L14	ll1 and ((gamma adj2 glutamyltranferase) or \$glutamyltransferase or (alpha adj2 glutamyltraspeptidase) or \$glutamyltranspeptidase or (gamma adj2 glutamyl adj (peptidyltransferase or transpeptidase)) or (\$glutamyl adj (peptidyltranferase or transpeptidase)) or (gamma adj glutamyltransferase) or \$glutamyltrasferase or (gamma adj gpt) or \$gpt or (gamma adj gt) or \$gt or (gamma adj gtp) or \$gt)	333
Γ.,	L13	L12 and l11	31
Г	L12	((diabetic adj neuropath\$) or glomeruloscleros\$ or glomerulopath\$ or ((chronic or persistent) adj (kidney or renal) adj disease))	7335
Г	L11	((At adj 125) or ACV or acivicin or (isoxazoleacetic adj acid))	1993
	DB=I	PGPB; PLUR=YES; OP=OR	
Γ.	L10	L8 and ((diabetic adj neuropath\$) or glomeruloscleros\$ or glomerulopath\$ or ((chronic or persistent) adj (kidney or renal) adj disease))	(
	DB=0	USPT; PLUR=YES; OP=OR	
Γ	L9	L8 and ((diabetic adj neuropath\$) or glomeruloscleros\$ or glomerulopath\$ or ((chronic or persistent) adj (kidney or renal) adj disease))	19
Г	L8	17 or 14	315
Γ	L7	\$isoxazoleacetic adj acid	∙ 2€
Γ	L6	L4 and (kidney or renal)	131
ľ.	L5	L4 and kidney or renal	23661
Γ	L4	12 or acivicin	294
Г	L3	L2.clm	131
Г	L2	At adj 125	131
Γ	L1	4758551.pn.	1

This is G o o g I e's cache of http://radiology.uchc.edu/eAtlas/GU/1505.htm as retrieved on Jun 18, 2005 10:35:33 GMT. Google's cache is the snapshot that we took of the page as we crawled the web.

The page may have changed since that time. Click here for the current page without highlighting.

This cached page may reference images which are no longer available. Click here for the cached text only.


To link to or bookmark this page, use the following url: http://www.google.com/search?

q=cache:LNB2V9DVV1EJ:radiology.uchc.edu/eAtlas/GU/1505.htm+etiology+%2B+focal+glomerulosclerosis&hl=en

Google is not affiliated with the authors of this page nor responsible for its content.

These search terms have been highlighted: etiology focal glomerulosclerosis

PathWeb Home Search Frames Search No frames Feed Back



Focal Segmental Glomerulosclerosis Click on Image to Enlarge it

#### Focal Segmental Glomerulosclerosis

- One segment of one glomerulus is sclerosed (arrow).
- •The other glomeruli are unremarkable.
- •With the trichrome stain, collagenous tissues take on a blue color.

(Description By:H. Yamase, M.D.) (Image Contrib. by: UCHC)

## **Focal Segmental Glomerulosclerosis**

# Etiology

- •Focal segmental glomerulosclerosis is not one disease but a histologic pattern of injury seen on light microscopy in patients with nephrotic range proteinuria.
- •Focal segmental glomerulosclerosis may be idiopathic or may be seen as a consequence of reflux nephropathy, AIDS nephropathy; heroin/drug addiction, or compensatory hypertrophy/hyperfiltration.

## **Pathogenesis**

•Since focal segmental glomerulosclerosis is not a disease so much as a histologic consequence of several diseases processes, the initiating pathogenetic mechanisms are varied. The common element to cases of focal segmental glomerulosclerosis is injury manifested by sclerosis in the setting of nephrotic range proteinuria.,

# **Epidemiology**

- •Focal segmental glomerulosclerosis is not one disease but a histologic pattern of injury seen on light microscopy in patients with nephrotic range proteinuria.
- •Focal segmental glomerulosclerosis may be idiopathic or may be seen as a consequence of reflux nephropathy, AIDS nephropathy; heroin/drug addiction, or compensatory hypertrophy/hyperfiltration.
- Immunogenetics: DR4.

## General Gross Description

- •The kidneys in focal segmental glomerulosclerosis would show no gross abnormalities in the early phases of the disease.
- •As the disease progresses and more glomeruli become sclerosed, the parenchymal mass will decrease and the kidneys would appear small and shrunken.

#### **General Microscopic Description**

- •The histologic recognition of the disease is dependent on examining a number of glomeruli together.
- Some of the glomeruli are normal and some show segmental sclerotic lesions.
- •The sclerotic lesions show solidification of segments of the tuft and is composed of collapsed capillary basement membranes, increased collagenous matrix and few if any viable endothelial and mesangial cells.
- •Some of the affected segments may show large hyaline deposits occluding previously patent capillary lumens.
- Direct immunofluorescence studies show IgM and C3 in the affected sclerosed segments.
- •Electron microscopy shows collapsed basement membranes, excess mesangial matrix material and the hyaline deposits noted on light microscopy correspond to large electron dense material occupying the capillary luminal space, which may represent entrapped serum proteins.
- Occasional foam cells may be encountered in the sclerosed segments.
- •The tubular compartment will show tracts of atrophic tubules, paralleling the degree of glomerulosclerosis.

#### Clinical Correlation

- •Patients with focal segmental glomerulosclerosis manifest heavy proteinuria (3.5 gm or more per 24 hours) and the consequence of protein loss (hypoalbuminemia, hyperlipidemia, edema).
- Additional complications of heavy proteinuria include increased vulnerability to infections, increased risk of thrombotic and thromboembolic diseases.
- Depending on the stage of the disease, patients may show normal renal function, varying degrees of renal insufficiency, or renal failure.

#### References

- Cotran RS, Kumar V, Robbins SL: Robbins Pathologic Basis of Disease. 5th ed. Philadelphia, W.B. Saunders, 1994, pp. 952-954.
- •Rose B. Renal Pathophysiology the essentials. Baltimore: Williams and Wilkins. 1994. Ch.

#### **Focal Segmental Glomerulosclerosis**

Synopsis by: Harold Yamase M.D. (T71200M00003)[212]

Search Medline at National Library of Medicine

Please be patient during transfer. Medline will open in a new window. To return, close the Medline Window

Search No frames PathWeb Home Feed Back Search Frames About

# ChemPort In-house Document Request

Send a copy of the reference below to your In-house Document esource.

io: STIC-ILL@uspto.gov

Contact Information: Change

JENNIFER HARLE 5712722763 jennifer.harle@uspto.gov

#### comments:

Please obtain the following article for SN 10/644325 and deliver to Remsen 3C18. i am in AU 1654 Room 3D28.

PLEASE NOTE THAT I NEED THE MONTH AND DATE THAT THIS ARTICLE WAS FIRST AVAILABLE. PLEASE OBTAIN THIS INFORMATION FOR ME AND LET ME KNOW WHEN YOU WILL HAVE IT AVAILABLE.

Send

#### **Document Citation**

**itle:** Transmembrane proteases as disease markers and targets for therapy.

**Authors:** Antczak C; de Meester I; Bauvois B

Source: J.Biol.Regul.Homeostatic Agents (15, No. 2, 130-39, 2001) 1 Fig. 3 Tab. 101 Ref. CODEN: JBRAER

SSN: 0393-974X

Corporate Source: INSERM; Inst.Curie; CNRS; Univ.Antwerp